

Research Article

Proximate Analysis on Ungingered and Gingered Tiger Nut's Drink Commercialized in Some Major Towns in Rivers State, Nigeria

Received:
February 24, 2023

Accepted:
March 5, 2023

Published:
March 12, 2023

Oghenerukevwe Obadesagbo^{1*}, Michael Horsefall Jnr¹, Lebari Sibe¹

¹Department of Biochemistry/Chemistry, Technology, School of Science Laboratory
Technology, University of Port Harcourt, Choba Rivers State, Nigeria.

How to cite:

Obadesagbo et al.
2023. "Proximate
Analysis on
Ungingered and
Gingered Tiger Nut's
Drink Commercialized
in Some Major Towns
in Rivers State,
Nigeria". Current
Research in
Interdisciplinary
Studies 2(3): 1 – 9.
<https://doi.org/10.5861/4/cris231>

Abstract: Fresh tigernut drink (*Cyperus esculentus*) was investigated for its proximate composition for ungingered and gingered tigernuts drink in some commercialized towns in Rivers State, Nigeria using the various methods described by the Association of Official Analytical Chemist (AOAC), 2010. The proximate analysis of the fresh tigernut drinks revealed that moisture content (86.51 ± 2.10 ; 82.26 ± 1.36) has the highest value followed by carbohydrate (7.20 ± 2.05 ; 10.17 ± 1.82), crude protein (2.83 ± 0.25 ; 3.34 ± 0.42) crude fat (1.62 ± 0.17 ; 2.50 ± 0.23), crude fiber (1.52 ± 0.18 ; 1.43 ± 0.31) and ash content (0.320 ± 0.16 ; 0.30 ± 0.03) respectively for ungingered and gingered tigernuts drinks. This study observed statistically significant variation ($P < 0.05$) in the moisture content, protein value, fat concentration and carbohydrate between the ungingered and gingered tigernut drinks. However, the fibre and ash contents showed no significant statistical variations ($P > 0.05$). In conclusion, the results provide additional information about the nutritional value of ungingered and gingered tigernuts drinks. The gingered tigernut drinks were revealed to contained higher level of energy-giving nutrients which is found to be beneficial for consumer's population in Port Harcourt metropolis.


Keywords: Proximate, Tigernut, Ungingered, Gingered, Nutrients.

Corresponding author:

Oghenerukevwe Obadesagbo
obadesagboangel@gmail.com

1. Introduction

Tigernut, or (*Cyperus esculentus*,) is a monocotyledonous perennial that grows to a maximum height of 24-55 cm and has a fibrous, upright root system (Maduka and Ire, 2018). The Cyperaceae family of plants is an excellent source of many nutrients, including but not limited to: energy, fat, carbs, fibre, glucose, and protein (Muhammed et al. 2011). Many essential nutrients, including vitamins and minerals, and digestive enzymes like catalase, lipase, and amylase, are abundant (Adejuyitan, 2011). Raw tiger-nut has been shown to contain several phytochemicals and antinutrients, including alkaloids, cyanogenic glycosides, resins, tannins, sterols, oxalates, phytates, and saponins (Chukwuma et al. 2010). Europe, South America, Asia, and Africa all have tigernut plantations. Food shortages have made this sweet tuber a popular dish (Bilikis and Olanrewaju, 2015). This was found to be the case by (Djomdi et al., 2013). Due to their availability throughout the year, tigernut tubers are routinely processed into new forms for consumption. In Africa, where tubers are widely consumed, this method of preparation is commonplace. These diets are inadequate for encouraging infant growth in developing countries due to their high anti-nutrient

 © 2023
by the authors. The
terms and conditions
of the Creative
Commons Attribution
(CC BY) licence apply
to this open access
article

content (Nnam 1995; Onilude et al. 1999). They have a better nutritional profile when mixed with other foods, which is well proven. It's also common knowledge that certain types of processing can increase a food's nutritional content by making it more digestible and reducing the amount of antinutritional chemicals (Ejoh et al. 2006). Many cultivars are produced, but only the yellow and brown varieties with some therapeutic characteristics make it to market (Suleiman et al. 2018; Farre, 2003). In particular, the yellow variety was in high demand because of its eye-catching colour, greater size, and meatier nuts. It produces more milk each lactation, is higher in protein content, and is lower in fat and anti-nutritional factors like polyphenols (Maduka and Ire, 2018). Such findings were published in 2013 (Musa and Hamza). Tigernut juice, a naturally energizing drink, is produced by blending crushed tigernut tubers with cinnamon, sugar, and vanilla. After ingesting this beverage, many people report feeling energised (Sánchez-Zapata et al., 2012).

Because of its high nutrient content, including carbs, glucose, proteins, minerals like phosphorus and potassium, and vitamins C and E, tigernut drink is a healthy and energising beverage. Similarly, it contains a high concentration of oleic acid, which strengthens the immune system and reduces the risk of contracting diseases including diarrhoea, constipation, and heart disease (Suleiman et al. 2018; Bixquert, 2003). Widespread consumption of tigernut drink in Nigeria has been linked to no adverse events (Onuoha et al. 2017). Several studies have been conducted on the nutritional composition of tigernut tuber variations from different locales (Nwaoguikpe, 2010). Nutritionists have taken an interest in tigernut tubers because of their widespread application in food production (Maduka and Ire, 2018). For more than just their ability to relieve thirst, people have been interested in and buying plant-based milk alternatives like tigernut in recent years. Because of its perishable nature, tiger-nut drink is best consumed within two to four hours at temperatures between forty and one hundred degrees Celsius (Akoma et al. 2006). Scientists discovered that tiger-nut milk, when pasteurised and flavoured with citric acid, ginger, and garlic, can stay fresh for an additional 2–3 days (Nwobosi et al. 2013). The anti-nutrient properties of the tigernut drink were drastically altered when spices were added as natural preservatives; these spices prevented the growth of micro-organisms throughout cold storage (from 5 days to indefinitely) (Kayode et al. 2021; Maxwell et al. 2018). This study aims to do just that by contrasting the nutritional profiles of ginger-infused and ginger-free tiger nut drinks already on the market.

2. Materials and Methods

2.1. Sample Collection

The tiger nut drink samples used in this study were purchased from three (3) vendors from different location in Choba, Eleme and Alakahia in pairs, stored with an ice chest container and then transported to the laboratory for proximate analysis.


2.2. Standards and Reagents

All reagents were of analytical reagent grade. De-ionized water was used for all dilutions. Nitric acid was not less than 65%, with a density of approximately 1.4 g/mL. Sulfuric acid (concentrated, N-free), Petroleum ether or Diethyl ether,

Catalyst/salt mixture (Kjeldahl digestion tablets), Nitrogen-free, anhydrous Sodium Sulphate, Sodium hydroxide solution, 50 %, w/v, Boric acid solution and Standardized HCl solution.

2.3. Sample preparation

Figure 1. Flow chart for Tigernut milk drink production



Source: Udeozor, (2012)

Figure 2. Picture of Tigernut seed and Milk extract



2.4. Sample analysis

Proximate analysis of ungingered and gingered tiger nuts drinks tigernuts

The standard methods were used to determine the dry matter, crude protein, total ash, crude fibre, and carbohydrate contents (AOAC, 2010). Water content was determined using the oven drying method at 105 °C for 5 hours, crude protein was measured using the micro-Kjeldahl method, total ash was obtained by igniting 2 g of sample in a muffle furnace at 550 °C for 4 hours, and crude fibre was measured using the digestion method to arrive at an estimate of carbohydrate content. Diethyl ether was employed in a standard soxhlet extraction to remove the fat (AOAC, 2010).

Quality assurance and quality control

All of the water used in the study was deionized, distilled twice, and purified (Prajapati and Patel 2010). The glassware and plasticware (Merck, Germany) were prewashed in 10% HNO₃, then rewashed in distilled water. The quality control results can be trusted because they were triple-checked using each sample. We followed the AOAC's suggested methods for proximity analysis to ensure the highest level of precision. The ICP Standard was used to develop certified reference solutions for individual elements, which were then used to calibrate the equipment (Merck, Germany).

2.5. Statistical Analysis

The laboratory data was analysed using IBM SPSS (version 23). One-way analysis of variance (ANOVA) was carried out to determine the significance level. Significance of differences was evaluated using a confidence interval spanning from 5% to 95%. Microsoft Excel and Word were used for the statistical and mathematical work. The data was presented as a Mean Standard Deviation.


3. Result and Discussion

Table 1. Proximate Analysis of Tiger Nut Drinks (*Cyperus esculentus*) from Port Harcourt Metropolis

Proximate Analysis	Ungingered Tiger Nut Drink (Mg/l)	Gingered Tiger Nut Drink (Mg/l)	F	Significance of Variation
Moisture	86.51 ± 2.10	82.26 ± 1.36	14.43	0.005
Fat	1.62 ± 0.17	2.50 ± 0.23	48.28	0.000
Protein	2.83 ± 0.25	3.34 ± 0.42	5.37	0.049
Fibre	1.52 ± 0.18	1.43 ± 0.31	0.31	0.593
Ash	0.320 ± 0.16	0.30 ± 0.03	0.09	0.773
Carbohydrate	7.20 ± 2.05	10.17 ± 1.82	5.88	0.042

Data are expressed as Mean ± Standard deviation

Figure 3. Variation in proximate analysis of Gingered and Ungingered Tiger nut drink from Port Harcourt Metropolis



Discussion

The mean concentration and standard deviation of the proximate composition of Gingered and Ungingered tiger nut (*Cyperus esculentus*) drink is presented in Table 1 while Figure 1 showed the variation in proximate analysis of gingered and ungingered tiger nut drink from Port Harcourt metropolis. The moisture content for the ungingered and gingered tiger nut drink was 86.51% and 82.26% respectively. These results were higher with those reported by Aremu *et al.* (2015); Oladele and Aina, (2007). The high moisture content may make the tiger nut drink relatively vulnerable to microbial attack. However, lower moisture content in gingered flavoured tiger nut could however reduce the rate of inhibition of the growth of microorganisms. Also, result from this research were also found to be higher when compared with moisture content values from treated and untreated unmalted tigernuts and treated and untreated malted tigernuts (Ndubuisi, 2015).

The fat content of these tiger nuts is lower than that found in the scholarly work of Ndubuisi, (2015) on the evaluation of food potentials of tigernut tubers (*Cyperus esculentus*) and its products (milk, coffee, and wine), but it is comparable to that of nuts and seeds, higher than that of cereals, and on par with soya beans (Sanchez-Zapata et al. 2012).

When compared to studies on the nutritional value of tiger nut (*Cyperus esculentus*) tubers and its products by Sabah et al. (2019) and studies on the nutritional value of protein-rich foods like soyabean, cowpeas, kersting nut, pigeon peas, and some oil seeds, the protein values for ungingered (2.83) and gingered (3.34) tiger nut drink are low (Oloafe et al. 2006).

The fibre content may improve digestion by lowering stomach pressure and speeding up the digestive process (Temple et al. 1990). As fibre helps reduce gas, it makes

sense to consider incorporating tiger nut fibre into meal plans for people struggling with digestive issues, constipation, and chronic conditions like colon cancer, diverticulitis, heart disease, and obesity (Wardlaw et al. 2002). The body relies heavily on crude fibre. It lengthens the amount of time waste takes to pass through the digestive system and increases stool volume. Most of the plant polysaccharides that make up crude fibre in the diet are indigestible by human digestive enzymes; they include cellulose, hemicellulose, and other components of plant cell walls (Southland, 1975). The fibre content of raw tiger nut, as reported by Oladele et al. (6.26% for the yellow variation and 5.62 for the brown variant), is significantly higher than the crude fibre values for ungingered and gingered tiger nut drinks (1.36% and 1.73%, respectively) (2007).

The ash concentration of both the ginger-infused and non-gingered beverages was very low, at between 0.11 and 0.56 percent, compared to the 1.18 percent found in tigernut (Suleiman et al. 2018).

Gingered tiger nut drink (10.17%) had a larger carbohydrate value than ungingered tiger nut drink (7.20%), making the former a more nourishing choice for energy generation. These findings exceeded those found in *Moringa oleifera*-flavored tiger nut drinks (3.600.06%) reported by Oyetoro et al. (2019). Moisture, carbohydrates, proteins, fats, fibres, and ashes were found to be present in descending order in the tiger nut drink proximate composition results. This means that the fat, fibre, and ash levels of tiger nut drinks are rather low.

Significant levels of proxies between ungingered and gingered Port Harcourt city tiger nuts were determined using one-way analysis of variance. Differences in moisture content, protein value, lipid concentration, and carbohydrate content between ungingered and ginger-infused tiger nuts were shown to be statistically significant ($P < 0.05$). There were no statistically significant differences in the levels of fibre or ash content, however ($P > 0.05$). Although all of the samples were generally accepted in terms of sensory quality, the research confirmed what Udeozor (2012) found when he analysed the proximal and sensory quality potentials of tiger nut-soy milk drink: a significant difference ($P < 0.05$) in taste and overall acceptability. Because of its high nutrient contents (protein, fat, etc.), it was determined that tiger-nut drink should be promoted in order to address the issue of protein-calorie malnutrition.

4. Conclusion

The results of this study provide new information about the nutritional value of tiger nut drinks with and without ginger. The research showed that the tigernuts drink used often in the Greater Port Harcourt Region is a nutritious and energising beverage due to its high protein content and abundance of carbs and lipids. The addition of different spices to tigernut drink such as ginger can increase the nutrients present in tigernut drink. The gingered tigernut drinks were observed to contained higher level of nutrients which is beneficial for adults and children population in Port Harcourt metropolis.

Acknowledgements

Many thanks go out to Mr. Ighariemu Victor of the Environmental Toxicology Units at the Department of Biochemistry in the Faculty of Science at the University of Port Harcourt.

Funding

There was no designated funding for this study from any governmental, commercial, or charitable institution.

Data Availability

If you ask the author of record, you can get data that backs up this work.

Declarations

Conflict of Interest

The people who wrote this article and anyone in their immediate families do not have any financial or other ties that could be seen as having influenced the work described here.

Consent for Publication

Each author has looked over the final draught and given it their approval.

References

- [1] Adejuyitan, J. A. 2011. "Tigernut processing: its food uses and health benefits". *American Journal of Food Technology*, 6(3), 197-201.
- [2] Akoma, O., Jiya, E. A., Akumka, D. D. & Mshelia, E. 2006. "Influence of malting on the nutritional characteristics of Kunun Zaki". *African Journal of Biotechnology*, 10(5): 996–1000.
- [3] Aremu M. O, Ibrahim H, Bamidele T.O, Agere H, Aremu SO. (2015). "Proximate composition and Amino acid profile of raw and cooked black variety of tiger nut (*Cyperus esculentus* L.) grown in Northeast Nigeria". *Journal of Biology, Agriculture and Healthcare* ISSN (5)7. 2221-3208
- [4] Association of Official Analytical Chemist (AOAC). *Official methods of analysis* (18th sed.). Washington DC (2010).
- [5] Bilikis, A., & Olanrewaju, A. 2015. "Chemical compositions, antioxidant capacity of tigernut (*Cyperus esculentus*) and potential health benefits". *European Scientific Journal*.
- [6] Bixquert, M. 2003. "Digestive aspects of tigernuts". *Pesca Y Alimentacion* 71, 85.
- [7] Chukwuma, E. R., Obioma, N., & Christopher, O. I. 2010. "The phytochemical composition and some biochemical effects of Nigerian tigernut (*Cyperus esculentus* L.) tuber". *Pakistan Journal of Nutrition*, 9(7), 709-715.
- [8] Djomdi, D., Kramer, J. K. G., VanderJagt, D. J., Ejoh, R., Ndjouenkeu, R., & Glew, R. H. 2013. "Influence of soaking on biochemical components of tiger nut (*Cyperus esculentus*) tubers cultivated in Cameroon". *International Journal of Food Process Engineering*, 1, 16-28.
- [9] Ejoh, R. A., Djomdi, & NDJOUENKEU, R. 2006. "Characteristics of tigernut (*Cyperus esculentus*) tubers and their performance in the production of a milky drink". *Journal of Food Processing and Preservation*, 30(2), 145-163.
- [10] Farre, R. 2003. "Nutritional and dietetic aspects of tiger nuts". *Food Science and Technology International* 11, 33-40.
- [11] <http://repository.futminna.edu.ng:8080/jspui/handle/123456789/10525>

- [12] Kayode, R. M., Joseph, J. K., Adegunwa, M. O., Dauda, A. O., Akeem, S. A., Kayode, B. I., ... & Olabanji, S. O. 2021. "Effects of addition of different spices on the quality attributes of tiger-nut milk (kunun-aya) during storage". *Journal of Microbiology, Biotechnology and Food Sciences*, 1-6.
- [13] Maduka, N., & Ire, F. S. 2018. "Tigernut plant and useful application of tigernut tubers (*Cyperus esculentus*) - A review". *Current Journal of Applied Science and Technology* 29(3), 1-23.
- [14] Maduka, N., & Ire, F. S. 2019. "A Review of Some Prevention Strategies against Contamination of *Cyperus esculentus* and Tigernut Derived Products of Economic Importance". *Asian Journal of Advanced Research and Reports* 1, 1-13.
- [15] Maxwell, Y. M. O., Alabi, M. O., & Edohe, A. O. 2018. "Effect of Spices on Anti-nutritional Properties of Tiger-nut Beverage (kunun aya)". *Proceedings of the 42nd Annual Conference and General Meeting of the Nigerian Institute of Food Science and Technology, Abeokuta, Nigeria*.
- [16] Muhammad, N. O., Bamishaiye, E. I., Bamishaiye, O. M., Usman, L. A., Salawu, M. O., Nafiu, M. O. & Oloyede, O. B. 2011. "Physicochemical Properties and Fatty Acid Composition of *Cyperus esculentus* (Tiger Nut) Tuber Oil". *Bioresearch Bulletin*, 5: 51- 54.
- [17] Musa, A. A., & Hamza, A. 2013. "Comparative Analysis of Locally Prepared 'Kunun Aya'(Tiger-Nut Milk) Consumed By Students Of Kaduna State University, Kaduna-Nigeria". *Science World Journal*, 8(2), 13-18.
- [18] Ndubuisi, L. C. 2015. "Evaluation of food potentials of tigernut tubers (*Cyperus esculentus*) and its products (milk, coffee and wine)" (Doctoral dissertation).
- [19] Nnam, N. M. 1995. "Evaluation of nutritional quality of fermented cowpea (*Vigna unguiculata*) flours". *Ecology of food and nutrition*, 33(4), 273-279.
- [20] Nwaoguikpe, R. N. 2010. "The phytochemical, proximate and amino acid compositions of the extracts of two varieties of tiger nut (*Cyperus esculentus*) and their effects on sickle cell hemoglobin polymerization. *Journal of Medical Sciences*, 1(11), 543-549.
- [21] Nwobosi, P. N. U., Isu, N. R. & Agarr, O. O. 2013. "Influence of pasteurization and use of natural tropical preservatives on the quality attributes of tiger nut drink during storage". *International Journal of Food and Nutrition Science*, 2: 27-32.
- [22] Oladele, A. K., & Aina, J. O. 2007. "Chemical composition and functional properties of flour from two varieties of tigernut". *African Journal Biotechnology*, 6, 2473 – 2476.
- [23] Olaofe, O., Akintayo, E. T., Adeyeye, E. I., & Adubiaro, H. O. 2006. "Proximate composition and functional properties of bulma cotton (*Bombacopsis glabra*) seeds". *Egypt Journal of Food Science*, 34, 81 – 90.
- [24] Onilude, A. A., Sanni, A. I., & Ighalo, M. I. 1999. "Effect of process improvement on the physico-chemical properties of infant weaning food from fermented composite blends of cereal and soybeans". *Plant Foods for Human Nutrition*, 54(3), 239-250.
- [25] Onuoha, N. O., Ogbusua, N. O., Okorie, A. N., & Ejike, C. E. 2017. "Tigernut (*Cyperus esculentus* L.) "milk" as a potent "nutri-drink" for the prevention of acetaminophen-induced hepatotoxicity in a murine model". *Journal of intercultural ethnopharmacology*, 6(3), 290.
- [26] Oyetoro AOA, Ogundipe OO, Adeyeye SAO, Akande EA, Akinyele AB. 2019. "Production and evaluation of tigernut (*Cyperus esculentus*) milk flavored with moringa oleifera leaf extract". *Current Research in Nutrition and Food Science* 7: 265–271.
- [27] Sabah, M. S., Shaker, M. A., Abbas, M. S., & Moursy, F. I. 2019. Nutritional value of Tiger nut (*Cyperus esculentus* L.) Tubers and its products. *Journal of Biological, Chemical and Environmental Sciences*, 14(1), 301-318.
- [28] Sánchez-Zapata, E., Fernández-López, J., & Angel Pérez-Alvarez, J. 2012. Tiger nut (*Cyperus esculentus*) commercialization: health aspects, composition, properties, and food applications. *Comprehensive Reviews in Food Science and Food Safety*, 11(4), 366-377.
- [29] Southland, W. M. 1975. "Biochemistry of Nutrition". Church Hill Livingstone, New York, pp. 471 – 473.

- [30] Suleiman, M. S., Olajide, J. E., Omale, J. A., Abbah, O. C., & Ejembi, D. O. 2018. "Proximate composition, mineral and some vitamin contents of tigernut (*Cyperus esculentus*)". *Clinical Investigation*, 8(4), 161-165.
- [31] Temple, V. J., Ojobe, T. O., & Kapu, M. M. 1990. Chemical analysis of tiger nut (*Cyperus esculentus*). *Journal of the Science of Food and Agriculture*, 50(2), 261-263.
- [32] Wardlaw GM, Kessel WM. 2002. "Calcium in food". *Perspective in nutrition* (5th ed.) 410-411.